

Cytokines

Bibliography of One Hundred Key Papers

-
- Agnoletti L, Curello S, Bachetti T, et al.** *Serum from patients with severe heart failure downregulates eNOS and is proapoptotic: role of tumor necrosis factor-alpha.*
Circulation. 1999;100:1983-1991.
-
- Anker SD, Clark AL, Kemp M, et al.** *Tumor necrosis factor and steroid metabolism in chronic heart failure: possible relation to muscle wasting.*
J Am Coll Cardiol. 1997;30:997-1001.
-
- Anker SD, Egerer KR, Volk HD, et al.** *Elevated soluble CD14 receptors and altered cytokines in chronic heart failure.*
Am J Cardiol. 1997;79:1426-1429.
-
- Anker SD, Volterrani M, Egerer KR, et al.** *TNF- α as predictor of peak leg blood flow in chronic heart failure.*
Q J Med. 1998;91:199-203.
-
- Aukrust P, Ueland T, Lien E, et al.** *Cytokine network in congestive heart failure secondary to ischemic or idiopathic dilated cardiomyopathy.*
Am J Cardiol. 1999;83:376-382.
-
- Bachetti T, Comini L, Agnoletti L, et al.** *Attivazione e ruolo del fattore di necrosi tumorale alfa nello scompenso cardiaco congestizio.*
Cardiologia. 1996;41:343-347.
-
- Balligand JL, Ungureanu D, Kelly RA, et al.** *Abnormal contractile function due to induction of nitric oxide synthesis in rat cardiac myocytes follows exposure to activated macrophage-conditioned medium.*
J Clin Invest. 1993;91:2314-2319.
-
- Bozkurt B, Kribbs S, Clubb FJ Jr, et al.** *Pathophysiologically relevant concentrations of tumor necrosis factor- α promote progressive left ventricular dysfunction and remodeling in rats.*
Circulation. 1998;97:1382-1391.
-
- Bozkurt B, Shan K, Seta Y, Oral H, Mann DL.** *Tumor necrosis factor- α and tumor necrosis factor receptors in human heart failure.*
Heart Fail Rev. 1996;1:211-219.
-
- Cain BS, Meldrum DR, Dinarello CA, et al.** *Tumor necrosis factor-alpha and interleukin-1 beta synergistically depress human myocardial function.*
Crit Care Med. 1999;27:1309-1318.
-
- Cecconi C, Curello S, Bachetti T, Corti A, Ferrari R.** *Tumor necrosis factor in congestive heart failure: a mechanism of disease for the new millennium?*
Prog Cardiovasc Dis. 1998;41(suppl 1):25-30.
-
- Chandrasekar B, Colston JT, Freeman GL.** *Induction of proinflammatory cytokine and antioxidant enzyme gene expression following brief myocardial ischaemia.*
Clin Exp Immunol. 1997;108:346-351.
-

Bibliography of One Hundred Key Papers

- Colucci WS.** *Molecular and cellular mechanisms of myocardial failure.*
Am J Cardiol. 1997;80:15L-25L.
-
- Deng MC, Erren M, Kammerling L, et al.** *The relation of interleukin-6, tumor necrosis factor-alpha, IL-2, and IL-2 receptor levels to cellular rejection, allograft dysfunction, and clinical events early after cardiac transplantation.*
Transplantation. 1995;60:1118-1124.
-
- Deswal A, Bozkurt B, Seta Y, et al.** *A phase I trial of tumor necrosis factor receptor (p75) fusion protein (TNFR:Fc) in patients with advanced heart failure.*
Circulation. 1999;99:3224-3226.
-
- Dibbs Z, Thornby J, White BG, Mann DL.** *Natural variability of circulating levels of cytokines and cytokine receptors in patients with heart failure: implications for clinical trials.*
J Am Coll Cardiol. 1999;33:1935-1942.
-
- Ferrari R, Bachetti T, Confortini R, et al.** *Tumor necrosis factor soluble receptors in patients with various degrees of congestive failure.*
Circulation. 1995;92:1479-1486.
-
- Ferrari R, Corti A, Bachetti T.** *Tumor necrosis factor alpha in heart failure.*
Heart Fail. 1995;(August/September):142-148.
-
- Finkel MS.** *Effects of proinflammatory cytokines on the contractility of mammalian heart.*
Heart Fail Rev. 1996;1:203-210.
-
- Finkel MS, Hoffman RA, Shen L, Oddis CV, Simmons RL, Hattler BG.** *Interleukin-6 (IL-6) as a mediator of stunned myocardium.*
Am J Cardiol. 1993;71:1231-1232.
-
- Finkel MS, Oddis CV, Jacob TD, Watkins SC, Hattler BG, Simmons RL.** *Negative inotropic effects of cytokines on the heart mediated by nitric oxide.*
Science. 1992;257:387-389.
-
- Frangogiannis NG, Lindsey ML, Michael LH, et al.** *Resident cardiac mast cells degranulate and release preformed TNF-alpha, initiating the cytokine cascade in experimental canine myocardial ischemia/reperfusion.*
Circulation. 1998;98:699-710.
-
- Givertz MM, Colucci WS.** *New targets for heart-failure therapy: endothelin, inflammatory cytokines, and oxidative stress.*
Lancet. 1998;352(suppl 1):S134-S138.
-
- Gulick T, Chung MK, Pieper SJ, Lange LG, Schreiner GF.** *Interleukin 1 and tumor necrosis factor inhibit cardiac myocyte beta-adrenergic responsiveness.*
Proc Natl Acad Sci USA. 1989;86:6753-6757.
-
- Gwechenberger M, Mendoza LH, Youker KA, et al.** *Cardiac myocytes produce interleukin-6 in culture and in viable border zone of reperfused infarctions.*
Circulation. 1999;99:546-551.
-
- Han RO, Ray PE, Baughman KL, Feldman AM.** *Detection of interleukin and interleukin-receptor mRNA in human heart by polymerase chain reaction.*
Biochem Biophys Res Commun. 1991;181:520-523.
-



-
- Hasper D, Hummel M, Docke WD, et al.** *Inflammatory mediators in patients with biventricular assist device systems.*
Z Kardiol. 1996;85:820-827.
-
- Hattler BG, Zeevi A, Oddis CV, Finkel MS.** *Cytokine induction during cardiac surgery: Analysis of TNF- α expression pre- and postcardiopulmonary bypass.*
J Card Surg. 1995;10:418-422.
-
- Hirota H, Chen J, Betz UA, et al.** *Loss of a gp130 cardiac muscle cell survival pathway is a critical event in the onset of heart failure during biomechanical stress.*
Cell. 1999;97:189-198.
-
- Hirota H, Yoshida K, Kishimoto T, Taga T.** *Continuous activation of gp130, a signal-transducing receptor component for interleukin 6-related cytokines, causes myocardial hypertrophy in mice.*
Proc Natl Acad Sci USA. 1995;92:4862-4866.
-
- Iwasaki A, Matsumori A, Yamada T, Shioi T, Wang W, Ono K, et al.** *Pimobendan inhibits the production of proinflammatory cytokines and gene expression of inducible nitric oxide synthase in a murine model of viral myocarditis.*
J Am Coll Cardiol. 1999;33:1400-1407.
-
- Kapadia SR.** *Cytokines and heart failure.*
Cardiol Rev. 1999;7:196-206.
-
- Kapadia S, Dibbs Z, Kurrelmeyer K, et al.** *The role of cytokines in the failing human heart. In: Crawford M, ed. Cardiology Clinics.* Philadelphia, Pa: W.B. Saunders; 1998:645-656.
-
- Kapadia S, Lee JR, Torre-Amione G, Birdsall HH, Ma TS, Mann DL.** *Tumor necrosis factor gene and protein expression in adult feline myocardium after endotoxin administration.*
J Clin Invest. 1995;96:1042-1052.
-
- Kapadia S, Oral H, Lee J, Nakano M, Taffet GE, Mann DL.** *Hemodynamic regulation of tumor necrosis factor- α gene and protein expression in adult feline myocardium.*
Circ Res. 1997;81:187-195.
-
- Kapadia S, Torre-Amione G, Mann DL.** *Pitfalls in measuring cytokines.*
Ann Intern Med. 1994;121:149-150.
-
- Kapadia S, Torre-Amione G, Yokoyama T, Mann DL.** *Soluble tumor necrosis factor binding proteins modulate the negative inotropic effects of TNF- α in vitro.*
Am J Physiol. 1995;37:H517-H525.
-
- Katz SD, Rao R, Berman JW, et al.** *Pathophysiological correlates of increased serum tumor necrosis factor in patients with congestive heart failure: relation to nitric oxide-dependent vasodilation in the forearm circulation.*
Circulation. 1994; 90:12-16.
-
- Kelly RA, Ballingand JL, Smith TW.** *Nitric oxide and cardiac function.*
Circ Res. 1996;79:363-380.
-
- Kelly RA, Smith TA.** *Cytokines and cardiac contractile function.*
Circulation. 1997; 95:778-781.
-
- Kishimoto T, Akira S, Narazaki M, Taga T.** *Interleukin-6 family of cytokines and gp130.*
J Am Soc Hematology. 1995;86:1243-1254.
-

Bibliography of One Hundred Key Papers

- Koller-Strametz J, Pacher R, Frey B, Kos T, Woloszczuk W, Stanek B.** *Circulating tumor necrosis factor-alpha levels in chronic heart failure: relation to its soluble receptor II, interleukin-6, and neurohumoral variables.*
J Heart Lung Transplant. 1998;17:356-362.
-
- Kubota T, Bounoutas GS, Miyagishima M, et al.** *Soluble tumor necrosis factor receptor abrogates myocardial inflammation, but not hypertrophy in cytokine-induced cardiomyopathy.*
Circulation. 2000;101:2518-2525.
-
- Kubota T, McNamara DM, Wang JJ, et al.** *Effects of tumor necrosis factor gene polymorphisms on patients with congestive heart failure.*
Circulation. 1998;97:2499-2501.
-
- Kubota T, McTiernan CF, Frye CS, Demetris AJ, Feldman AM.** *Cardiac-specific overexpression of tumor necrosis factor-alpha causes lethal myocarditis in transgenic mice.*
J Card Fail. 1997;3:117-124.
-
- Kubota T, McTiernan CF, Frye CS, et al.** *Dilated cardiomyopathy in transgenic mice with cardiac specific overexpression of tumor necrosis factor-alpha.*
Circ Res. 1997;81:627-635.
-
- Levine B, Kalman J, Mayer L, Fillit HM, Packer M.** *Elevated circulating levels of tumor necrosis factor in severe chronic heart failure.*
N Engl J Med. 1990;223:236-241.
-
- Li YY, Feldman AM, Sun Y, McTiernan CF.** *Differential expression of tissue inhibitors of metalloproteinases in the failing human heart.*
Circulation. 1998;98:1728-1734.
-
- Li YY, McTiernan CF, Feldman AM.** *Proinflammatory cytokines regulate tissue inhibitors of metalloproteinases and disintegrin metalloproteinase in cardiac cells.*
Cardiovasc Res. 1999;42:162-172.
-
- MacGowan GA, Kormos RL, McNamara DM, Alvarez RJ, Rosenblum WD, Pham S, et al.** *Predicting short-term outcome in severely ill heart failure patients: implications regarding listing for urgent cardiac transplantation and patient selection for temporary ventricular assist device support.*
J Card Fail. 1998;4:169-175.
-
- MacGowan GA, Mann DL, Kormas RL, Feldman AM, Murali S.** *Circulating interleukin-6 in severe heart failure.*
Am J Cardiol. 1997;79:1128-1131.
-
- Mann DL.** *Stress activated cytokines and the heart.*
Cytokine Growth Factor Rev. 1996;7:341-354.
-
- Mann DL.** *Cytokines as mediators of disease progression in the failing heart. In: Hosenpud JD, Greenberg BH, eds.*
Congestive Heart Failure. Philadelphia, Pa: Lippincott-Williams & Wilkins; 1999:213-232.
-
- Mann DL.** *Inflammatory mediators in heart failure: homogeneity through heterogeneity.*
Lancet. 1999; 353:1812-1813.
-
- Mann DL, Lee-Jackson D, Yokoyama T.** *Tumor necrosis factor-alpha and cardiac remodeling.*
Heart Fail. 1995;(September/October):166-175.
-



- Mann DL, Young JB.** *Basic mechanisms in congestive heart failure: recognizing the role of proinflammatory cytokines.*
Chest. 1994;105:897-904.
-
- Matsui S, Matsumori A, Matoba Y, Uchida A, Sasayama S.** *Treatment of virus-induced myocardial injury with a novel immunomodulating agent, vesnarinone.*
J Clin Invest. 1994;94:1212-1217.
-
- Matsumori A, Ono K, Nishio R, Nose Y, Sasayama S.** *Amiodarone inhibits production of tumor necrosis factor- α by human mononuclear cells. A possible mechanism for its effect in heart failure.*
Circulation. 1997;96:1386-1389.
-
- Matsumori A, Shioi T, Yamada T, Matsui S, Sasayama S.** *Vesnarinone, a new inotropic agent, inhibits cytokine production by stimulated human blood from patients with heart failure.*
Circulation. 1994;89:955-958.
-
- Matsumori A, Yamada T, Kawai C.** *Immunomodulating therapy in viral myocarditis: effects of tumour necrosis factor, interleukin 2 and anti-interleukin-2 receptor antibody in an animal model.*
Eur Heart J. 1991;12:203-205.
-
- Matsumori A, Yamada T, Matoba Y, Sasayama S.** *Tumor necrosis factor in cardiomyopathy and myocarditis.*
Circulation. 1992;86(suppl I):1439. Abstract.
-
- Matsumori A, Yamada T, Suzuki H, Matoba Y, Sasayama S.** *Increased circulating cytokines in patients with myocarditis and cardiomyopathy.*
Hypertension. 1989;13:706-711.
-
- McTiernan CF, Lemster BH, Frye C, Brooks S, Combes A, Feldman AM.** *Interleukin-1 β inhibits phospholamban gene expression in cultured cardiomyocytes.*
Circ Res. 1997;81:493-503.
-
- Meldrum DR.** *Tumor necrosis factor in the heart.*
Am J Physiol. 1998;274:R577-R595.
-
- Meldrum DR, Cain BS, Cleveland JC Jr, et al.** *Adenosine decreases post-ischemic cardiac TNF- α production: anti-inflammatory implications for preconditioning and transplantation.*
Immunology. 1997;92:472-477.
-
- Muller-Werdan U, Engelmann H, Werdan K.** *Cardiodepression by tumor necrosis factor-alpha.*
Eur Cytokine Netw. 1998;9:689-691.
-
- Muller-Werdan U, Schumann H, Loppnow H, et al.** *Endotoxin and tumor necrosis factor alpha exert a similar proinflammatory effect in neonatal rat cardiomyocytes, but have different cardiodepressant profiles.*
J Mol Cell Cardiol. 1998;30:1027-1036.
-
- Munger MA, Johnson B, Amber IJ, Callahan KS, Gilbert EM.** *Circulating concentrations of proinflammatory cytokines in mild or moderate heart failure secondary to ischemic or idiopathic dilated cardiomyopathy.*
Am J Cardiol. 1996;77:723-727.
-
- Nakano M, Knowlton AA, Dibbs Z, Mann DL.** *Tumor necrosis factor- α confers resistance to injury induced by hypoxic injury in the adult mammalian cardiac myocyte.*
Circulation. 1998;97:1392-1400.
-
- Nakano M, Knowlton AA, Yokoyama T, Lesslauer W.** *Tumor necrosis factor- α induced expression of heat shock protein 72*

Bibliography of One Hundred Key Papers

- Mann DL.** *in adult feline cardiac myocytes.*
Am J Physiol. 1996;270:H1231-H1239.
-
- Niebauer J, Pflaum CD, Clark AL, et al.** *Deficient insulin-like growth factor I in chronic heart failure predicts altered body composition, anabolic deficiency, cytokine and neurohormonal activation.*
J Am Coll Cardiol. 1998;32:393-397.
-
- Niebauer J, Volk HD, Kemp M, et al.** *Endotoxin and immune activation in chronic heart failure: a prospective cohort study.*
Lancet. 1999;353:1838-1842.
-
- Oddis CV, Finkel MS.** *Cytokine-stimulated nitric oxide production inhibits mitochondrial activity in cardiac myocytes.*
Biochem Biophys Res Commun. 1995;213:1002-1009.
-
- Oddis CV, Simmons RL, Hattler BG, Finkel MS.** *Chronotropic effects of cytokines and the nitric oxide synthase inhibitor, L-NMMA, on cardiac myocytes.*
Biochem Biophys Res Commun. 1994;205:992-997.
-
- Oddis CV, Simmons RL, Hattler BG, Finkel MS.** *cAMP enhances inducible nitric oxide synthase mRNA stability in cardiac myocytes.*
Am J Physiol. 1995;269:H2044-H2050.
-
- Ono K, Matsumori A, Shioi T, Furukawa Y, Sasayama S.** *Cytokine gene expression after myocardial infarction in rat hearts: possible implication in left ventricular remodeling.*
Circulation. 1998;98:149-156.
-
- Oral H, Dorn GW, II, Mann DL.** *Sphingosine mediates the immediate negative inotropic effects of tumor necrosis factor- α in the adult mammalian cardiac myocyte.*
J Biol Chem. 1997;272:4836-4842.
-
- Oral H, Kapadia S, Nakano M, et al.** *Tumor necrosis factor- α and the failing human heart.*
Clin Cardiol. 1995;18(suppl IV):IV20-IV27.
-
- Packer M.** *Is tumor necrosis factor an important neurohormonal mechanism in chronic heart failure?*
Circulation. 1995;92:1379-1382.
-
- Paulus WJ.** *How are cytokines activated in heart failure?*
Eur J Heart Fail. 1999;1:309-312.
-
- Pulkki KJ.** *Cytokines and cardiomyocyte death.*
Ann Med. 1997;29:339-343.
-
- Sato Y, Matsumori A, Sasayama S.** *Inotropic agent vesnarinone inhibits cytokine production and E-selectin expression in human umbilical vein endothelial cells.*
J Mol Cell Cardiol. 1995;27:2265-2273.
-
- Schreiner GF.** *Immune modulation of cardiac cell function.*
Trans Am Clin Climatol Assoc. 1998;109:39-49; discussion 49-50.
-
- Seta Y, Shan K, Bozkurt B, Oral H, Mann DL.** *Basic mechanisms in heart failure: the cytokine hypothesis.*



-
- J Card Fail.* 1996;2:243-249.
-
- Shan K, Kurrelmeyer K, Seta Y, et al.** *The role of cytokines in disease progression in heart failure.*
Curr Opin Cardiol. 1997;12:218-223.
-
- Sheng Z, Knowlton K, Chen J, Hoshijima M, Brown JH, Chien KR.** *Cardiotrophin 1 (CT-1) inhibition of cardiac myocyte apoptosis via a mitogen-activated protein kinase-dependent pathway.*
J Biol Chem. 1997;272:5783-5791.
-
- Shimokawa H, Kuroiwa-Matsumoto M, Takeshita A.** *Cytokine generation capacities of monocytes are reduced in patients with severe heart failure.*
Am Heart J. 1998;136:991-1002.
-
- Shioi T, Matsumori A, Sasayama S.** *Persistent expression of cytokine in the chronic stage of viral myocarditis in mice.*
Circulation. 1996;94:2930-2937.
-
- Sliwa K, Skudicky D, Candy G, Wisenbaugh T, Sareli P.** *Randomized investigation of effects of pentoxifylline on left ventricular performance in idiopathic dilated cardiomyopathy.*
Lancet. 1998;351:1091-1093.
-
- Testa M, Yeh M, Lee P, et al.** *Circulating levels of cytokines and their endogenous modulators in patients with mild to severe congestive heart failure due to coronary artery disease or hypertension.*
J Am Coll Cardiol. 1996;28:964-971.
-
- Torre-Amione G, Kapadia S, Benedict CR, Oral H, Young JB, Mann DL.** *Proinflammatory cytokine levels in patients with depressed left left ventricular ejection fraction: a report from the studies of left ventricular dysfunction (SOLVD).*
J Am Coll Cardiol. 1996;27:1201-1206.
-
- Torre-Amione G, Kapadia S, Lee J, Bies RD, Lebovitz R, Mann DL.** *Expression and functional significance of tumor necrosis factor receptors in human myocardium.*
Circulation. 1995;92:1487-1493.
-
- Torre-Amione G, Kapadia S, Lee J, et al.** *Tumor necrosis factor- α and tumor necrosis factor receptors in the failing human heart.*
Circulation. 1996;93:704-711.
-
- Torre-Amione G, MacLellan WR, Kapadia S, et al.** *Tumor necrosis factor- α is persistently expressed in cardiac allografts in the absence of histological or clinical evidence of rejection.*
Transplant Proc. 1998;30:875-877.
-
- Torre-Amione G, Stetson SJ, Youker KA, et al.** *Decreased expression of tumor necrosis factor-alpha in failing human myocardium after mechanical circulatory support: a potential mechanism for cardiac recovery.*
Circulation. 1999;100:1189-1193.
-
- Tsutamoto T, Hisanaga T, Wada A, et al.** *Interleukin-6 spillover in the peripheral circulation increases with the severity of heart failure, and the high plasma level of interleukin-6 is an important prognostic predictor in patients with congestive heart failure.*
J Am Coll Cardiol. 1998;31:391-398.
-
- Turner D, Grant SC, Yonan N, et al.** *Cytokine gene polymorphism and heart transplant rejection.*

Bibliography of One Hundred Key Papers

- Ungunreanu-Longrois D, Balligand JL, Okada I, et al.** *Contractile responsiveness of ventricular myocytes to isoproterenol is regulated by induction of nitric oxide synthase activity in cardiac microvascular endothelial cells in heterotypic primary culture.* **Circ Res.** 1995;77:486-493.
-
- Vanderheyden M, Kersschot E, Paulus WJ.** *Pro-inflammatory cytokines and endothelium-dependent vasodilation in the forearm. Serial assessment in patients with congestive heart failure.* **Eur Heart J.** 1998;19:747-752.
-
- Wagner DR, Combes A, McTiernan CF, Sanders VJ, Lemster B, Feldman AM.** *Adenosine inhibits lipopolysaccharide-induced cardiac expression of tumor necrosis factor- α .* **Circ Res.** 1998;82:47-56.
-
- Wagner DR, Kubota T, Sanders VJ, McTiernan CF, Feldman AM.** *Differential regulation of cardiac expression of IL-6 and TNF- α by A₂- and A₃-adenosine receptors.* **Am J Physiol.** 1999;276(6 pt 2):H2141-H2147.
-
- Wagner DR, McTiernan CF, Sanders VJ, Feldman AM.** *Adenosine inhibits lipopolysaccharide-induced secretion of tumor necrosis factor- α in the failing human heart.* **Circulation.** 1998;97:521-524.
-
- Wollert KC, Taga T, Saito M, et al.** *Cardiotrophin-1 activates a distinct form of cardiac muscle cell hypertrophy.* **J Biol Chem.** 1996;271:9535-9545.
-
- Yamauchi-Takahara K, Ihara Y, Ogata A, Yoshizaki K, Azuma J, Kishimoto T.** *Hypoxic stress induces cardiac myocyte-derived interleukin-6.* **Circulation.** 1995;91:1520-1524.
-
- Yokoyama T, Nakano M, Bednarczyk JL, McIntyre BW, Entman ML, Mann DL.** *Tumor necrosis factor- α provokes a hypertrophic growth response in adult cardiac myocytes.* **Circulation.** 1997;95:1247-1252.
-
- Yokoyama T, Vaca L, Rossen RD, Durante W, Hazarika P, Mann DL.** *Cellular basis for the negative inotropic effects of tumor necrosis factor- α in the adult mammalian heart.* **J Clin Invest.** 1993;92:2303-2312.
-
- Yoshida K, Taga T, Saito M, et al.** *Targeted disruption of gp130, a common signal transducer for the interleukin 6 family of cytokines, leads to myocardial and hematological disorders.* **Proc Natl Acad Sci USA.** 1996;93:407-411.
-
- Yue P, Massie BM, Simpson PC, Long CS.** *Cytokine expression increases in nonmyocytes from rats with postinfarction heart failure.* **Am J Physiol.** 1998;275:H250-H258.
-