

Stroke

Bibliography of One Hundred Key Papers

selected by **Giuseppe Mancina, MD** and **Guido Grassi, MD**
Clinica Medica – Ospedale S. Gerardo – Monza (Milano) - ITALY

-
- Algra A, van Gijn J.** *Aspirin at any dose above 30 mg offers only modest protection after cerebral ischaemia.*
J Neurol Neurosurg Psychiatry. 1996;60:197-199.
-
- ALLHAT Collaborative Research Group.** *Major cardiovascular events in hypertensive patients randomized to doxazosin vs chlorthalidone.*
JAMA. 2000;283:1967-1975.
-
- Alter M, Sobal E, McCoy R, et al.** *Stroke in the Lehigh Valley: risk factors for recurrent stroke.*
Neurology. 1987;37:503-507.
-
- American Heart Association.** *AHA Practice guidelines for the use of imaging in transient ischaemic attacks and acute stroke.*
Stroke. 1997;28:1480-1497.
-
- Antiplatelet Trialists' Collaboration.** *Secondary prevention of vascular disease by prolonged antiplatelet treatment.*
BMJ. 1988;296:320-331.
-
- Bamford J, Sandercock P, Dennis M, et al.** *A prospective study of acute cerebrovascular disease in the community: the Oxfordshire Community Stroke Project—1981-1986: methodology, demography and incident cases of first-ever stroke.*
J Neurol Neurosurg Psychiatry. 1988;51:1373-1380.
-
- Barnett H, Taylor D, Eliasziw H, et al.** *Benefit of carotid endarterectomy in patients with symptomatic moderate or severe stenosis.*
N Engl J Med. 1998;339:1415-1425.
-
- Blauw GJ, Lagaay AM, Smelt AH, et al.** *Stroke, statins and cholesterol: a meta-analysis of randomized, placebo-controlled, double-blind trials with HMG-CoA reductase inhibitors.*
Stroke. 1997;28:946-950.
-
- Blood Pressure Lowering Treatment Trialists' Collaboration.** *Effects of ACE-inhibitors, calcium antagonists and other blood-pressure-lowering drugs: results of prospectively designed overview of randomized trials.*
Lancet. 2000;356:1955-1964.
-
- Bonita R.** *Epidemiology of stroke*
Lancet. 1992;339:342-344.
-
- Borhani NO, Mercuri M, Borhani PA, et al.** *Final outcome results of the Multicenter Isradipine Diuretic Atherosclerosis Study (MIDAS). A randomized controlled trial.*
JAMA. 1996;276:785-791.
-



-
- Bots ML, Hoes AW, Koudstaal PJ, Hofman A, Grobbee DE.** *Common carotid intima-media thickness and risk of stroke and myocardial infarction—the Rotterdam study.*
Circulation. 1997;96:1432-1437.
-
- Brott T, Bogousslavsky J.** *Treatment of acute ischaemic stroke.*
N Engl J Med. 2000;343:710-722.
-
- Burn J, Tennis M, Banford J, Sandercock P, Wade D, Warlow C.** *Long-term risk of recurrent stroke after a first-ever stroke: the Oxfordshire Community Stroke Project.*
Stroke. 1994;25:1605-1610.
-
- CAPRIE Steering Committee.** *A randomized, blinded, trial of clopidogrel versus aspirin in patients at risk of ischaemic events (CAPRIE).*
Lancet. 1996;348:1329-1339.
-
- Collins R, MacMahon S.** *Blood pressure, antihypertensive drug treatment and risk of stroke and of coronary heart disease.*
Br Med Bull. 1994;50:272-298.
-
- Dahlof B, Devereux RB, Kjeldsen SE, et al.** *Cardiovascular morbidity and mortality in the Losartan Intervention For the Endpoint reduction in hypertension study (LIFE).*
Lancet. 2002;359:995-1003.
-
- Diener HC, Cunha L, Forbes C, Silvenius J, Smets P, Lowenthal A.** *European Stroke Prevention study 2: dipyridamole and acetylsalicylic acid in the secondary prevention of stroke.*
J Neurol Sci. 1996;143:1-13.
-
- Estacio RO, Jeffers BW, Hiatt WR, et al.** *The effects of nisoldipine as compared with enalapril on cardiovascular outcomes in patients with non-insulin dependent diabetes and hypertension.*
N Engl J Med. 1998;338:645-652.
-
- European Carotid Surgery Trialists' Collaborative Group.** *Randomized trial of endarterectomy for recently symptomatic carotid stenosis: final results of the MRC European Carotid Surgery Trial (ECST).*
Lancet. 1998;351:1379-1387.
-
- Forette F, Seux ML, Staessen JA, et al.** *Prevention of dementia in randomized double-blind placebo-controlled Systolic Hypertension in Europe (Syst-Eur) Trial.*
Lancet. 1998;352:1347-1351.
-
- Gent M, Blakely JA, Easton JD, et al.** *The Canadian American Ticlopidine Study (CATS) in thromboembolic stroke.*
Lancet. 1989;1:1215-1220.
-
- Gillum LA, Mamidipudi SK, Johnston SC.** *Ischaemic stroke risk with oral contraceptives: a meta-analysis.*
JAMA. 2000;284:72-78.
-
- Goldstein LB, Adams R, Becker K, et al.** *Primary prevention of ischaemic stroke.*
Stroke. 2001;32:280-299.
-
- Golledge J, Greenhalgh RM, Davies AH.** *The symptomatic carotid plaque.*
Stroke. 2000;31:774-781.
-
- Gueyffier F, Boissel JP, Boutitie F, et al.** *Effect of antihypertensive treatment in patients having already suffered from stroke. Gathering the evidence.*
Stroke. 1997;28:2557-2562.
-

Bibliography of One Hundred Key Papers

- Guidelines Subcommittee 1999 World Health Organization / International Society of Hypertension.** *Guidelines for the management of hypertension.* **J Hypertens.** 1999;17:151-183.
-
- Hacke W, Kaste M, Fieschi C, et al, for the ECASS Study Group.** *Intravenous thrombolysis with recombinant tissue plasminogen activator for acute hemispheric stroke: the European Cooperative Acute Stroke Study (ECASS).* **JAMA.** 1995;274:1017-1025.
-
- Hademenos GJ, Massoud TF.** *Biophysical mechanisms of stroke.* **Stroke.** 1997;28:2067-2077.
-
- Hankey GJ, Warlow CP.** *Treatment and secondary prevention of stroke: evidence, costs and effects on individual populations.* **Lancet.** 1999;354:1457-1463.
-
- Hansson L, Zanchetti A, Carruthers SG, et al.** *Effects of intensive blood pressure lowering and low-dose aspirin in patients with hypertension: principal results of the Hypertension Optimal Treatment (HOT) randomised trial.* **Lancet.** 1998;351:1755-1762.
-
- Hass WK, Easton JD, Adams HP, et al.** *A randomized trial comparing ticlopidine hydrochloride with aspirin for the prevention of stroke in high-risk patients: Ticlopidine Aspirin Stroke Study Group.* **N Engl J Med.** 1989;321:501-507.
-
- Heart Outcomes Prevention Evaluation Study Investigators.** *Effects of an angiotensin-converting enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients.* **N Engl J Med.** 2000;342:145-153.
-
- Heiss WD, Thiel A, Grond M, Graf R.** *Which targets are relevant for therapy of acute ischaemic stroke?* **Stroke.** 1999;30:1486-1489.
-
- Hellemons BSP, Langenberg M, Lodder J, et al.** *Primary prevention of arterial thromboembolism in non-rheumatic atrial fibrillation in primary care: randomized controlled trial comparing two intensities of coumarin with aspirin.* **BMJ.** 1999;319:958-964.
-
- Herbert PR, Graziano JH, Cham KS, Hennekens CH.** *Cholesterol lowering with statin drugs, risk of stroke, and total mortality. An overview of randomized trials.* **JAMA.** 1997;278:319-321.
-
- Hirsh J, Fuster V.** *Guide to anticoagulant therapy. Part 2: oral anticoagulants.* **Circulation.** 1994;89:1469-1480.
-
- Hobson RWI, Weiss DG, Fields WS, et al, for the VA Cooperative Study Group.** *Efficacy of carotid endarterectomy for asymptomatic carotid stenosis: the Veterans Affairs Cooperative Study Group.* **N Engl J Med.** 1993;328:221-227.
-
- Inzitari D, Eliasziw M, Gates P, et al.** *The causes and risk of stroke in patients with asymptomatic internal carotid artery stenosis.* **N Engl J Med.** 2000;342:1693-1700.
-
- Isles CG, Walzer LM, Bewers GD, et al.** *Mortality in patients of the Glasgow Blood Pressure Clinic.* **J Hypertens.** 1986;4:141-156.
-



-
- Jick H, Zornberg GL, Jick SS, Seshadri S, Drachman DA.** *Statins and the risk of dementia.*
Lancet. 2000;356:1627-1631.
-
- Johnston KC, Li JY, Lyden PD, Candhal S, et al.** *Medical and neurological complications of ischaemic stroke: experience from the RANTAS trial.*
Stroke. 1998;29:447-453.
-
- Joint National Committee on Prevention, Detection, Evaluation and Treatment of High Blood Pressure.** *The sixth report of the Joint National Committee on Prevention, detection, evaluation and treatment of high blood pressure. (JNC VI).*
J Hypertens. 2000;18:1795-1800.
-
- Kaira L, Evans A, Perez I, Knapp M, Donaldson N, Swift CG.** *Alternative strategies for stroke care: a prospective randomized controlled trial.*
Lancet. 2000;356:894-899.
-
- Kannel WB, McGee DL.** *Diabetes and cardiovascular disease: the Framingham Study.*
JAMA. 1979;241:2035-2038.
-
- Khaw KT.** *Epidemiology of stroke.*
J Neurol Neurosurg Psychiatry. 1996;61:333-338.
-
- Kjeldsen SE, Julius S, Hedner T, Hansson L.** *Stroke is more common than myocardial infarction in hypertension: analysis based on 11 major randomized intervention trials.*
Blood Pressure. 2001;10:190-192.
-
- Kiely DK, Wolf PA, Cupples LA, et al.** *Familial aggregation of stroke: the Framingham study.*
Stroke. 1993;24:1366-1371.
-
- Kitamura A, Iso H, Iida H, et al.** *Trends in incidence of coronary heart disease and stroke and the prevalence of cardiovascular risk factors among Japanese men from 1963 to 1994.*
Am J Med. 2002;112:104-109.
-
- Kjeldsen SE, Erdine S, Farsang C, Sleight P, Mancia G.** *1999 WHO/ISH Hypertension Guidelines: highlights and European Society of Hypertension update.*
J Hypertens. 2002;20:153-155.
-
- Liao DP, Cooper L, Cai JW, et al.** *Presence and severity of cerebral white matter lesions and hypertension, its treatment and its control: the ARIC Study.*
Stroke. 1996;27:2262-2270.
-
- Lip GY.** *Thromboprophylaxis for atrial fibrillation.*
Lancet. 1999;354:4-6.
-
- MacMahon S.** *Blood pressure and prevention of stroke.*
J Hypertens. 1996;14(suppl 6):539-546.
-
- MacMahon S, Peto R, Cutler J, et al.** *Blood pressure, stroke and coronary heart disease. Part I. Prolonged differences in blood pressure: prospective observational studies corrected for regression dilution bias.*
Lancet. 1990;335:765-774.
-
- MacMahon S, Rogers A.** *The epidemiological association between blood pressure and stroke: implications for primary and secondary prevention.*
Hypertens Res. 1994;17(suppl 1):S23-S32.
-

Bibliography of One Hundred Key Papers

- Mancia G, Giannattasio C, Grassi G.** *Current antihypertensive treatment: can we do better?*
Am J Hypertens. 1999;12:1315-1385.
-
- Mancia G, Grassi G.** *Antihypertensive treatment: past, present and future.*
J Hypertens. 1998;16(suppl 1):51-57.
-
- Mancia G, Grassi G.** *Rationale for the use of fixed combination in the treatment of hypertension.*
Eur Heart J. 1999;20(suppl 2):214-219.
-
- Mancia G, Grassi G.** *World Health Organization / International Society of Hypertension Guidelines and the role of ongoing trials.*
Eur Heart J. 2000;21(suppl D):D7-D10.
-
- Mancia G, Grassi G.** *Blood pressure reduction as part of the multifactorial approach to treatment of cardiovascular disease.*
Eur Heart J. 2001;22(suppl B):B32-B36.
-
- Mancia G, Groppelli A, Casadei R, Omboni S, Mutti E, Parati G.** *Cardiovascular effects of smoking.*
Clin Exp Hypertens. 1990;A12: 917-929.
-
- Marks MP, Marcellus M, Norbash AM, Steinberg GK, Tong DC, Albers GW.** *Outcome of angioplasty for atherosclerotic intracranial stenosis.*
Stroke. 1999;30:1065-1069.
-
- Okamoto K, Yamori Y, Nagaoka A.** *Establishment of the stroke-prone spontaneously hypertensive rat (SHR).*
Circ Res. 1974;34:1143-1153.
-
- Ott A, Breteler MMB, Van Harskamp F, Stijnen T, Hofman A.** *Incidence and risk of dementia: the Rotterdam Study.*
Am J Epidemiol. 1998;147:574-580.
-
- Park JB, Schiffrin EL.** *Effects of antihypertensive therapy on hypertensive vascular disease.*
Curr Hypertens Rep. 2000;2:280-288.
-
- PATS Collaborating Group.** *Post-stroke antihypertensive treatment. A preliminary result.*
Clin Med J. 1995;108:710-717.
-
- PROGRESS Collaborative Group.** *Randomized trial of a perindopril-based blood pressure-lowering regimen among 6105 individuals with previous stroke or transient ischaemic attack.*
Lancet. 2001;358:1033-1041.
-
- Qizilbash N, Lewington S, Duffy S, et al.** *Cholesterol, diastolic blood pressure and stroke. 13 000 strokes in 45 0000 people in 45 prospective cohorts: Prospective Studies Collaboration.*
Lancet. 1995;346:1647-1653.
-
- Sacco RL, Benjamin EJ, Broderick JP, et al.** *American Heart Association Prevention Conference IV. Prevention and rehabilitation of stroke. Risk factors.*
Stroke. 1997;28:1507-1517.
-
- Sacco RL, Boden-Albala B, Gan R, et al.** *Stroke incidence among white, black and Hispanic residents of an urban community: the Northern Manhattan Stroke Study.*
Am J Epidemiol. 1998;147:259-1268.
-
- Safar ME.** *Pulse pressure, arterial stiffness and cardiovascular risk.*
Curr Opin Cardiol. 2000;15:258-263.
-



-
- Safar ME, Blacher J, Mourad JJ, London GM.** *Stiffness of carotid artery wall material and blood pressure in humans. Application to antihypertensive therapy and stroke prevention.* **Stroke.** 2000;31:782-790.
-
- Scandinavian Simvastatin Survival Study Group.** *Randomized trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S).* **Lancet.** 1994;344:1383-1389.
-
- Sever PS, Dalhof B, Poulter NR, et al.** *Rationale, design, methods and baseline demography of participants of the Anglo-Scandinavian Cardiac Outcomes Trial.* **J Hypertens.** 2001;19:1139-1147.
-
- SHEP Cooperative Research Group.** *Prevention of stroke by antihypertensive drug treatment in older persons with isolated systolic hypertension: final results of the Systolic Hypertension in the Elderly Program (SHEP).* **JAMA.** 1991;265:3255-3264.
-
- Shinton R, Beevers G.** *Meta-analysis of relation between cigarette smoking and stroke.* **BMJ.** 1989;298:789-794.
-
- Skoog I, Nilsson L, Palmertz B, Andreasson LA, Svanborg A.** *A population-based study of dementia in 85-year-olds.* **N Engl J Med.** 1993;328:153-158.
-
- Stroke Prevention in Atrial Fibrillation Investigators.** *Warfarin versus aspirin for prevention of thromboembolism in atrial fibrillation: stroke prevention in atrial fibrillation II study.* **Lancet.** 1994;343:687-691.
-
- Stroke Unit Trialists' Collaboration.** *Collaborative systematic review of the randomized trials of organized inpatient (stroke unit) care after stroke.* **BMJ.** 1997;314:1151-1159.
-
- Sytkowski PA, D'Agostino RB, Belanger AJ, et al.** *Secular trends in long-term sustained hypertension, long-term treatment, and cardiovascular mortality; the Framingham Heart Study 1950-1990.* **Circulation.** 1996;93:697-703.
-
- Tatemichi TK, Paik M, Bagiella E, et al.** *Risk of dementia after stroke in a hospitalized cohort: results of a longitudinal study.* **Neurology.** 1999;44:1885-1891.
-
- The Diabetes Control and Complications Trial Research Group.** *The effect of intensive treatment of diabetes on the development and progression of long-term complication in insulin-dependent diabetes mellitus.* **N Engl J Med.** 1993;329:977-986.
-
- The INDIANA (Individual Data Analysis of Antihypertensive Intervention Trials) Project Collaborators.** *Effect of antihypertensive treatment in patients having already suffered from stroke: gathering the evidence.* **Stroke.** 1997;28:2557-2562.
-
- The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group.** *Tissue plasminogen activator for acute ischaemic stroke.* **N Engl J Med.** 1995;346:1509-1514.
-
- Thomson R, Parkin D, Eccles M, Sudlow M, Robinson A.** *Decision analysis and guidelines for anticoagulation therapy to prevent stroke in patients with atrial fibrillation.* **Lancet.** 2000;355:956-962.
-

Bibliography of One Hundred Key Papers

- Thorvaldsen P, Asplund K, Kuulasmaa K, Rajakangas AM, Schroll M, for the WHO MONICA project.** *Stroke incidence, case fatality and mortality in the WHO MONICA project.* **Stroke.** 1995;26:361-367.
-
- UKPDS Final Report.** *34th Annual Meeting of the European Association for the Study of Diabetes.* Barcelona, Spain, September 8-12, 1998. 1998.
-
- UK Prospective Diabetes Study (UKPDS) Group.** *Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type II diabetes (UKPDS 33).* **Lancet.** 1998;352:837-853.
-
- UK Prospective Diabetes Study Group.** *Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38.* **BMJ.** 1998;317:703-713.
-
- Vasan RS, Larson MG, Leip EP, et al.** *Impact of high-normal blood pressure on the risk of cardiovascular disease.* **N Engl J Med.** 2001;245: 1291-1297.
-
- Walker SP, Rimm EB, Ascherio A, et al.** *Body size and fat distribution as predictors of stroke among US men.* **Am J Epidemiol.** 1996;144:1143-1150.
-
- Wannamethee SG, Shaper AG, Whincup PH, et al.** *Smoking cessation and the risk of stroke in middle-aged men.* **JAMA.** 1995;274:155-160.
-
- White HD, Simes RJ, Anderson NE, et al.** *Pravastatin therapy and the risk of stroke.* **Engl J Med.** 2000;343:317-326.
-
- Wilhelmsen L.** *Epidemiology of smoking and intervention studies of smoking.* **Am Heart J.** 1988;115:242-249.
-
- Wolf PA, Abbott RD, Kannel WB.** *Atrial fibrillation as an independent risk factor of stroke: the Framingham Study.* **Stroke.** 1991;22:983-988.
-
- Wolf PA, D'Agostino RB, Belanger AJ, Kannel WB.** *Probability of stroke: a risk profile from the Framingham Study.* **Stroke.** 1991;22:312-318.
-
- Wolf PA, D'Agostino RB, O'Neal MA, et al.** *Secular trends in stroke incidence and mortality: the Framingham study.* **Stroke.** 1992;23:1551-1555.
-
- Wood D, De Backer G, Faergeman O, Graham I,** *Prevention of coronary heart disease in clinical practice. Recommendations of the Second Joint Task Force of European and other societies on coronary prevention.* **J Hypertens.** 1988;16:1407-1414.
-
- World Health Organization.** *The World Health Report 1999.* 1999. Geneva, Switzerland: WHO; 1999.
-